Four Pathways to Recovery

An expanded, printable diagrammatic guide for interrupting limerence

Purpose

This expanded guide breaks the four recovery pathways into practical actions. It is designed to be used alongside the 20-stage limerence pathway, especially when intrusive thoughts, compulsive contact, shame, or relationship risk remain active.

Recovery is not a single method. It can combine self-directed change, therapy, support from trusted people, values-based action, and attention to health and wellbeing.

Expanded recovery diagram

The diagram shows four pathways and explains how they work together. Use the actions that are most relevant to your current situation.

FOUR PATHWAYS TO RECOVERY

Four complementary routes can work together to interrupt limerence and rebuild a stable life.

PATHWAY 1■IMMEDIATE STABILISATION

- Reduce optional contact
- Make necessary contact professional, brief and neutral
- Remove digital reinforcement: no checking, rereading or monitoring
- Use grounding, sleep, exercise, food and routine after exposure
- Delay urges: wait 24 hours before acting on an impulse
- Plan what to do after unavoidable contact

PATHWAY 2■CHANGE THE THINKING

- Label limerent thoughts: “This is a thought, not a fact”
- Reality-test interpretations: separate evidence from assumption
- Challenge destiny beliefs: “I feel intensely” is not proof of fate
- Reduce rumination: postpone analysis, write once, then refocus
- Use CBT-style thought records and behavioural experiments
- Practise self-compassion rather than shame or self-attack

PATHWAY 3■REBUILD LIFE AND IDENTITY

- Invest directly in the marriage: communication, affection, repair
- Reconnect with values: partner, parent, colleague, friend, self
- Build support: trusted friends, family, community or peer support
- Create meaning beyond the LO: work, learning, hobbies, fitness
- Restore routines: sleep, meals, exercise, work and leisure
- Develop a future that does not depend on the LO’s attention

PATHWAY 4■SEEK HELP WHEN NEEDED

- Consider individual therapy for rumination, shame and boundaries
- Consider couples therapy if the marriage needs repair or support
- Use CBT for intrusive thoughts, compulsions and avoidance
- Use ACT to act according to values even when thoughts persist
- Seek urgent help for hopelessness, self-harm risk or loss of control
- Review progress regularly and adjust the plan as needed

HOW THE FOUR PATHWAYS WORK TOGETHER

Stabilise the situation → Change the thinking → Rebuild life and identity → Seek help when needed

The aim is not simply to stop thinking about the LO.

The aim is to build a life that is no longer organised around the LO’s attention, approval or absence.

If there is hopelessness, inability to stay safe, or thoughts of self-harm, seek urgent help immediately.

In Australia: Lifeline 13 11 14 • Emergency: 000

Figure 1. Four complementary pathways to recovery from limerence.

Pathways at a glance

The table summarises the purpose, actions, and main target of each pathway.

Pathway	Purpose	Core actions	What it interrupts	When to prioritise
1. Immediate stabilisation	Reduce exposure and emotional activation quickly.	Reduce optional contact; make necessary contact neutral; remove digital reinforcement; use grounding, sleep, exercise and routine.	The reward-seeking loop and compulsive checking.	When contact, urges, checking or emotional overwhelm are high.
2. Change the thinking	Reduce rumination, fantasy and destiny beliefs.	Label thoughts; reality-test interpretations; challenge “soul mate” beliefs; postpone analysis; use CBT or ACT strategies.	Rumination, idealisation and false hope.	When thoughts are intrusive, repetitive or emotionally consuming.
3. Rebuild life and identity	Restore sources of meaning beyond the LO.	Invest in marriage, family, values, friendships, work, health, hobbies and future goals.	Neglect of self and others; emotional dependence.	When life has become organised around the LO or the primary relationship needs repair.
4. Seek help when needed	Gain structured support and address underlying vulnerability.	Individual or couples therapy; CBT for compulsions and rumination; ACT for values-based action; urgent support for safety concerns.	Shame, isolation, repeated relapse and escalating risk.	When self-help is not enough, distress persists, or safety is a concern.

A practical weekly plan

Choose one action from each pathway. Small, repeated actions are more useful than relying on willpower alone.

Pathway	One action for this week	How I will know it is working
Immediate stabilisation	Remove one source of optional contact or checking.	Fewer urges to check, message or seek proximity.
Change the thinking	Label three limerent thoughts as thoughts, not facts.	Less time spent analysing ambiguous behaviour.
Rebuild life and identity	Schedule one protected activity with my partner, family or friend.	More connection and meaning outside the LO.
Seek help when needed	Book an assessment or therapy session, or tell one safe person.	Greater support, accountability and clearer boundaries.

Safety

If there is hopelessness, inability to stay safe, or thoughts of self-harm, seek urgent help immediately. In Australia, call Lifeline on **13 11 14** or **000** in an emergency.

Educational and reflective document. Not a substitute for assessment or treatment by a qualified mental-health professional.